# Testing hot star atmospheres with nebular modeling

Daniel Schaerer[1], Christophe Morisset[2], Jean-Claude Bouret[2], Fabrice Martins[1]

[1] *Observatoire Midi-Pyrénées, 14, Av. E. Belin, F-31400 Toulouse*

[1] *Laboratoire d'Astrophysique de Marseille, Traversé du Siphon-Les trois Lucs, F-13376 Marseille Cedex 12*

As the bulk of radiation in early type stars is emitted in the Lyman continuum and thus inaccessible to direct observations (except in extremely rare cases) it is crucial to find indirect tests to constrain this part of the spectrum, where the essential atmospheric conditions are settled. Nebular observations combined with detailed photoionization modeling can in principle provide such strongly needed constraints.

Here we present the current status of available constraints on the ionizing fluxes of O stars and first results from extensive photoionization model grids using recent non-LTE line blanketed model atmospheres including stellar winds and plane parallel models for O stars (CoStar: Schaerer & de Koter 1997, WMBASIC: Pauldrach et al. 2001, CMFGEN: Hillier & Miller 1998, TLUSTY: Hubeny & Lanz 2001).

The method and its dependence on nebular parameters (geometry etc.) are discussed. Our models are compared to a sample of HII regions observed with ISO providing a measure of the ionization degree through numerous IR fine structure lines. For stars with Teff $\leq$ 40000 K, all most recent atmosphere models show serious deficiencies in the wavelength range below $\sim$ 400 Angstrøm, predicting too weak fluxes. The possible causes for such a discrepancy are examined.