




Question 1: An Application of Larmor’s Formula: Thomson Scattering and the Thomson Cross Section



One of the simplest applications of Larmor’s formula is the scattering of radiation by a free electron. In the classical approximation, this process is called Thomson scattering. It is obtained by considering a sinusoidal electromagnetic wave with E -vector

$$E(t) = E_0 \sin \omega t \quad (\text{w1.1})$$

interacting with an electron at rest.

- a) Assuming $v \ll c$, show that the force on the electron is

$$F = m_e \ddot{\mathbf{r}} = e E_0 \sin \omega t \quad (\text{w1.2})$$

and that the dipole moment of the electron is given by

$$d = -\frac{e^2 E_0}{m_e \omega^2} \sin \omega t \quad (\text{w1.3})$$

- b) Using the results from the previous question and the dipole approximation, calculate the time averaged power emitted per unit angle and the total power emitted by the electron.

Reminder: $\langle \sin^2 \omega t \rangle = 1/2$.

- c) The *differential cross section*, $d\sigma/d\Omega$, is a measure how much radiation is scattered into a certain direction. It is defined through

$$\frac{dP}{d\Omega} = \langle S \rangle \frac{d\sigma}{d\Omega} \quad (\text{s1.1})$$

where the incident flux of radiation on the electron is given by Poynting’s theorem as

$$\langle S \rangle = \frac{c}{8\pi} E_0^2 \quad (\text{s1.2})$$

From your previous results show that

$$\frac{d\sigma}{d\Omega} = \frac{e^4}{m_e^2 c^4} \sin^2 \theta = r_0^2 \sin^2 \theta \quad (\text{s1.3})$$

where

$$r_0 = \frac{e^2}{m_e c^2} = 2.82 \times 10^{-13} \text{ cm} \quad (\text{s1.4})$$

is called the *classical electron radius*.

- d) The *total cross section* for the scattering of radiation off a free electron is obtained by integrating $d\sigma/d\Omega$ over Ω or immediately from

$$P = \langle S \rangle \sigma \quad (\text{s1.1})$$

Show that

$$\sigma = \frac{8\pi}{3} r_0^2 =: \sigma_T \quad (\text{s1.2})$$

where

$$\sigma_T = 6.652 \times 10^{-25} \text{ cm}^2 \quad (\text{s1.3})$$

is called the *Thomson cross section*.

Question 2: Parseval's Theorem

The Fourier transform pair as used in this lecture was defined by

$$F(\omega) = \int_{-\infty}^{+\infty} f(t)e^{i\omega t} dt \quad \text{and} \quad f(t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} F(\omega)e^{-i\omega t} d\omega \quad (\text{w2.1})$$

Provided that $f(t)$ is a sufficiently nice function, show that Parseval's theorem holds

$$\int |f(t)|^2 dt = \frac{1}{2\pi} \int |F(\omega)|^2 d\omega = \int |F(\nu)|^2 d\nu \quad (\text{w2.2})$$

Reminder 1: $|f|^2 = ff^*$ where f^* is the complex conjugate of f

Reminder 2: The δ -function can be written as

$$\delta(t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{i\omega t} d\omega \quad (4.34)$$